

Prediction of Ultrasonic Velocities in Binary Liquid Mixtures of N,N-Dimethyl Acetamide With Certain Amines

V. Sravanthi, P. Paul Divakar, K. Samatha

Abstract—Ultrasonic velocity and density values have been measured for the binary liquid mixtures of N,N Dimethyl Acetamide as a common component with Diethylamine, Diethanolamine and Benzylamine at 303, 308, 313 and 318K over the entire composition range. Theoretical velocity values have been evaluated by various relations viz., Nomoto, Free Length Theory (FLT), Van deal and Vangeel ideal mixing relation (IMR), Impedance Dependence Relation (IDR), and Junjie for three binary liquid mixtures. An attempt has been made to compare the merits of the relations and the relative applicability of these theories to the present systems have been checked and discussed. The results are explained in terms of intermolecular interactions occurring in these binary systems. The deviation in the variation of $U^2_{\text{exp}} / U^2_{\text{imx}}$ from unity has also been evaluated for explaining the non-ideality in the mixtures

Index Terms—ultrasonic velocity; binary liquid mixture; theoretical velocity; intermolecular interactions; non-ideality.

I. INTRODUCTION

Now a days much interest has been shown on the study of physico-chemical properties, behaviour and molecular interactions in liquid mixtures. The analysis of thermodynamic properties of the liquid mixtures can be used to get the qualitative information about the energetic and structural effects and packing phenomena that govern the mixing process. Ultrasonic velocity studies in binary and multi component liquid mixtures which are capable to reveal the hydrogen bonding among the molecules has been carried out by many researchers [1-3].

Theoretical evaluation of ultrasonic velocity in binary liquid mixtures and its correlation to study molecular interaction has been successfully done in recent years [4-6]. Ultrasonic velocities of liquid mixtures are of considerable importance in understanding intermolecular interaction between component molecules, and they find applications in several industrial and technological processes. Several relations, semi-empirical formula and theories are available for the theoretical computation of ultrasonic velocity in liquid and

liquid mixtures.

Vysyaraju Sravanthi, Department of Physics, Andhra University, Viskhapatnam, India,

Putla Paul Divakar, Department of Physics, Sir. C.R..ReddyAutonomous College, Eluru, India.

K. Samatha, Department of Physics, Andhra University, Viskhapatnam, India.

Using the theories available in literature, ultrasonic velocities in liquid mixtures have been calculated and compared with those obtained experimentally. The comparison of theoretical ultrasonic velocities with those obtained experimentally reveals the nature of interactions between the component molecules in the mixtures.

The aim of the present investigation is to compare the ultrasonic sound velocity in three binary liquid mixtures from various theoretical relations of Nomoto, Free Length Theory (FLT), Van deal and Vangeel ideal mixing relation (IMR), Impedance Dependence Relation (IDR), and Junjie. An attempt has been made to compare the merits of the relations for the binary liquid mixtures investigated at four different temperatures. The relative applicability of these theories to the present systems have been checked and discussed. The results are explained in terms of intermolecular interactions occurring in these binary systems. The deviation in the variation of $U^2_{\text{exp}} / U^2_{\text{imx}}$ from unity has also been evaluated for explaining the non-ideality of the mixtures.

II. EXPERIMENTAL

The binary liquid mixtures under study are:

- 1.N,N-Dimethyl Acetamide + Diethylamine (NNDA+DEA)
- 2.N,N-DimethylAcetamide + Diethanolamine (NNDA +DEIA) and
- 3.N,N-DimethylAcetamide + Benzylamine (NNDA +BA)

All the chemicals used in the present research work are AR grade of minimum assay of 99.9% and hence used without further purification. The liquid mixtures were prepared by mixing calculated amount of pure liquids. The ultrasonic velocities in the liquid mixtures were measured using a single crystal variable path interferometer operating at a frequency of 10MHz (MITTAL ENTERPRISES, New Delhi, Model:M-84) with an overall accuracy of $\pm 0.1\%$. The temperature during the experiment was controlled by circulating water around the liquid cell from the thermostatically controlled adequately stirred water bath. The densities of pure liquids and liquid mixtures were determined from the weight measurements using 10ml specific gravity bottle by the standard procedure with an accuracy of $\pm 0.1\text{kg m}^{-3}$.

III. THEORETICAL

3.1 NOMOTO'S RELATION:

Rao found experimentally that [7], for pure liquids, the ratio of temperature coefficients of sound velocity U and molar

volume remains almost constant:
$$\left(\frac{1}{U}\right)\left(\frac{dU}{dT}\right) = -3 \left(\frac{1}{V}\right)\left(\frac{dV}{dT}\right)$$

Where T is the absolute temperature. On Integrating this equation we get

$$VU^{1/3} = \text{const} \tan t = \frac{M}{\rho U^{1/3}} = R$$

Where M is molecular weight and ρ is density. The constant R is called the molar sound velocity or Rao’s constant. It was found to be additive i.e., it can be calculated as a sum of increments from the atoms or atom groups in the molecule and from the chemical bonds. On assuming the additivity of molar sound velocity (R) and no volume change on mixing, Nomoto established the following relation for a liquid mixture

$$R = \frac{M}{\rho U^{1/3}}$$

Where U and ρ are determined experimentally and M is the mean molecular weight in a binary liquid mixture

$$M = (X_1M_1 + X_2M_2)$$

Where X₁ and X₂ are the mole fractions and M₁ and M₂ are molecular weights of constituent components respectively. Simple theoretical treatment gives the following relation

$$U_{Nomoto} = \left[\frac{(X_1R_1 + X_2R_2)}{(X_1V_1 + X_2V_2)} \right]^3 \quad (1)$$

3.2 Relation based on Free Length Theory

Jacobson[8] deduce an empirical relation for ultrasonic velocity (U_{FLT}) making use of intermolecular free length (L_f) and density (ρ) as

$$U_{FLT} = \left[\frac{K}{L_{f,mix} \rho_{exp}^{1/2}} \right] \quad (2)$$

Where K is temperature dependent called Jacobson’s constant and the value of K at the working temperatures of the experiment were calculated (MKS units) and they are given below

| | | | | |
|-----------------------|-------------------|-------------------|-------------------|-------------------|
| Temp ⁰ (K) | 303 | 308 | 313 | 318 |
| Value of K | 2.075 | 2.095 | 2.115 | 2.135 |
| | ×10 ⁻⁶ | ×10 ⁻⁶ | ×10 ⁻⁶ | ×10 ⁻⁶ |

3.3 Ideal Mixing Relation based on the Van Deal and Vangeel Theory

Van Deal and Vangeel[9] proposed the ideal mixing theory in the light of assumptions made by Blandamer and Wadding[10], yield the following relation for adiabatic compressibility (β_{ad})_{mix}

$$(\beta_{ad})_{mix} = \Phi_1 \frac{\gamma_1}{\gamma_{mix}} (\beta_{ad})_1 + \Phi_2 \frac{\gamma_2}{\gamma_{mix}} (\beta_{ad})_2$$

Where Φ₁ and Φ₂ volume fractions of the liquids 1 and 2, γ₁ and γ₂ are the ratios of specific heats of the respective liquids. This relation holds good if the mixture is ideal and if γ₁=γ₂=γ_{mix}. Using the additional assumption that V₁=V₂ the above equation can be transformed into a linear combination of mole fraction X₁ and X₂,

$$(\beta_{ad})_{mix} = X_1 (\beta_{ad})_1 + X_2 (\beta_{ad})_2$$

On the basis of this equation, Van Deal and Vangeel obtained the relation for ultrasonic velocity in liquid mixtures as

$$U_{IMR} = \left[\frac{1}{X_1M_1 + X_2M_2} \right]^{1/2} \left[\frac{X_1}{M_1U_1^2} + \frac{X_2}{M_2U_2^2} \right]^{-1/2} \quad (3)$$

Where U₁ and U₂ are the ultrasonic velocities of the pure liquid components.

3.4 Impedance Dependence Relation

The ultrasonic velocity can be evaluated by the Impedance Dependence Relation [11] of the following form

$$U_{IDR} = \frac{X_1Z_1 + X_2Z_2}{X_1\rho_1 + X_2\rho_2} \quad (4)$$

where X₁ and X₂ are the mole fractions, ρ₁ and ρ₂ are the densities and Z₁ and Z₂ are the acoustic impedances of the liquid components.

3.5 Junjie’s Relation

Junjie’s Relation[12] for ultrasonic velocity is given by

$$U_{Junjie} = \left[\frac{X_1V_1 + X_2V_2}{(X_1M_1 + X_2M_2)^{1/2}} \right] \left[\frac{X_1V_1}{\rho_1U_1^2} + \frac{X_2V_2}{\rho_2U_2^2} \right]^{-1/2} \quad (5)$$

Where, 1 & 2 represents the first and second component of the liquid mixture and the other symbols have their usual meanings.

3.6 Percentage Deviation

The percentage deviation is calculated from the relation

$$\% \text{ deviation} = \sum \frac{(U_{mix(obs)} - U_{mix(cal)})}{U_{mix(obs)}} \times 100 \quad (6)$$

Here, U_{mix(obs)} is experimental value of ultrasonic velocity and U_{mix(cal)} is computed value of ultrasonic velocity. The worst-case error is the maximum value of deviation of the theoretical values from experimental values of ultrasonic velocity.

3.7 Degree of interaction, α

The deviation of the ratio U²_{exp} / U²_{imx} from unity is called degree of interaction, α.

IV. RESULTS AND DISCUSSION

The theoretical relations used to calculate ultrasonic velocity in all the above liquid mixtures are mentioned above at different temperatures 303, 308, 313 and 318K and the values along with the experimental values are given in the Tables from 1 to 3 Also the validity of different theoretical formulae is checked by percentage deviation for all the mixtures and at all the temperatures are shown.

It is observed from the tabulated values that the theoretical values of ultrasonic velocities evaluated by the above mentioned relations show deviations from the experimental values. The reason for the deviation maybe the limitations and approximations incorporated in these theories. The effect of volume change due to mixing was not considered in the Nomoto’s relation. That is interaction between the molecules was not taken into account. In Free Length theory, it was assumed that the molecules are of spherical shape but it is not true at all times. In the case of Ideal mixing relation, it was assumed that, the ratio of specific heats and volumes are equal. Again no molecular interactions were considered.

Upon mixing two liquids, the interaction between the molecules of the two liquids take place because of the

presence of various types of forces such as dispersion forces, charge transfer, hydrogen bonding, dipole-dipole and dipole-induced-dipole interactions. Thus, the observed deviation of theoretical values of velocity from the experimental values shows the molecular interactions between the unlike molecules in the liquid mixture [13-18]. In the present study, in the three binary mixtures esters, the velocities predicted by the Nomoto's relation are in better agreement than the other relations. When two liquids are mixed various types of forces play a vital role due to interactions. Thus, the observed deviations between theoretical and experimental values of velocity shows that there is molecular interaction between the unlike molecules in the liquid mixture. The FLT assumes that molecules are rigid spheres with no interaction between them and it is not valid in all the cases. Hence the deviation from the experimental ultrasonic velocity values is maximum in the FLT.

4.1 NN-Dimethylacetamide + Diethylamine

From the reported values, Table-1, it can be observed that the experimental values are very close in agreement with the computed values by Nomoto's relation, followed by the values obtained by Junjie's relation, Ideal Mixing Relation Impedance Dependence Relation and Free Length Theory. The percentage deviations are very less positive in the case of Nomoto's values, whereas the deviations are large for FLT relation with negative values at all temperatures. The Nomoto's Relation gives the best approximation for this binary mixture.

4.2 NN-Dimethylacetamide + Diethanolamine

A perusal of Table-2 reveals that the values calculated on the basis of Nomoto's Relation are in good agreement with the experimental values and there is no significant deviation at all temperatures.

Next to that, the values obtained by Nomoto's relation are very close to that of the experimental values and a slight negative deviation is observed. The values evaluated by the Junjie's relation, Ideal Mixing Relation and IDR follows

the above respectively with a little more deviations. For this mixture also the FLT gives a large deviation in evaluated ultrasonic velocity values compared to the values obtained by the other four relations.

4.3 NN-Dimethylacetamide + Benzylamine

A close look at Table-3 reveals that velocities determined by Nomoto's Relation give a very close approximation with the measured values and the deviation observed is almost nil.

The deviation in values calculated by Junjie's relation, Impedance Dependence Relation and Ideal Mixing Relation follows the deviation of the values obtained by respectively. In this case also the observed fact is that, FLT approximated values are much higher and negative values, the percentage deviations are also high.

The positive values of α in all the system clearly indicate the existence of tendency for the formation of association in mixture through hydrogen bonded complexes [17,18].

Velocities were determined on the basis of different theories and relations are discussed by other researchers earlier [19-21] and the validity of different theoretical formulae is checked by percentage deviation at different temperatures. As per the earlier studies, the limitations and approximations incorporated in these theories are responsible for the deviations between theoretical and experimental values.

Figs 1, 2 and 3 represent the variation of U^2_{exp} / U^2_{imx} with mole fraction of NN-Dimethylacetamide. It is observed that the curves are similar in all the three systems with maximum approximately at 0.5 mole fraction of NN-Dimethylacetamide at all temperatures. They are increasing with increase in temperature. The trend of the curves reveals a fact that the mixtures tend to move towards more non-ideality up to the middle mole fraction of NN-Dimethylacetamide, which suggests the formation of association in liquid mixtures through dipole-dipole interactions as reported by Shukla et al [22].

Graphical representation for variation of U^2_{exp} / U^2_{imx} with mole fraction of N,N Dimethyl acetamide




Fig.1 for NNDA+DEA




Fig.2 for NNDA +DEIA




Fig.3 for NNDA +BA

Table 1. Experimental and Theoretical values of Ultrasonic velocity and Percentage of Deviation at different temperatures for the binary liquid mixture- I (NN-Dimethylacetamide + Diethylamine).

| Mole Fraction of NNDA | Exp. Velocity | Theoretical Velocities | | | | | Percentage Deviation | | | | | $\frac{U^2}{U_{imx}^2}$ |
|-----------------------|---------------|------------------------|-----------|-----------|-----------|-----------|----------------------|------------|-----------|-----------|-----------|-------------------------|
| | | X_1 | U_{exp} | U_{Nom} | U_{FLT} | U_{IMR} | U_{IDR} | U_{JUNJ} | U_{Nom} | U_{FLT} | U_{IMR} | |
| 303K | | | | | | | | | | | | |
| 0.0000 | 1084.8 | 1084.8 | 1084.8 | 1084.8 | 1084.8 | 1084.8 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.0000 |
| 0.2149 | 1139.0 | 1154.0 | 997.2 | 1132.0 | 1176.5 | 1123.9 | 1.30 | -14.22 | -0.62 | 3.19 | -1.34 | 1.0124 |
| 0.3811 | 1189.0 | 1210.4 | 932.1 | 1177.1 | 1241.4 | 1163.3 | 1.77 | -27.56 | -1.01 | 4.22 | -2.21 | 1.0203 |
| 0.5135 | 1234.0 | 1257.3 | 892.8 | 1219.5 | 1289.7 | 1202.1 | 1.86 | -38.21 | -1.19 | 4.32 | -2.65 | 1.0239 |
| 0.6215 | 1274.0 | 1296.9 | 879.2 | 1259.1 | 1327.1 | 1240.0 | 1.77 | -44.91 | -1.18 | 4.00 | -2.74 | 1.0238 |
| 0.7112 | 1311.1 | 1330.7 | 883.7 | 1296.0 | 1356.9 | 1277.0 | 1.47 | -48.37 | -1.17 | 3.37 | -2.67 | 1.0235 |
| 0.7870 | 1345.2 | 1359.9 | 906.0 | 1330.3 | 1381.1 | 1312.8 | 1.08 | -48.48 | -1.12 | 2.60 | -2.47 | 1.0226 |
| 0.8518 | 1374.1 | 1385.4 | 964.5 | 1362.2 | 1401.3 | 1347.5 | 0.82 | -42.46 | -0.88 | 1.94 | -1.97 | 1.0176 |
| 0.9079 | 1400.2 | 1407.8 | 1057.7 | 1391.9 | 1418.3 | 1381.2 | 0.54 | -32.38 | -0.60 | 1.28 | -1.37 | 1.0120 |
| 0.9568 | 1424.1 | 1427.7 | 1200.8 | 1419.6 | 1432.9 | 1413.9 | 0.25 | -18.60 | -0.32 | 0.61 | -0.72 | 1.0064 |
| 1.0000 | 1445.5 | 1445.5 | 1445.5 | 1445.5 | 1445.5 | 1445.5 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.0000 |
| 308K | | | | | | | | | | | | |
| 0.0000 | 1069.8 | 1069.8 | 1069.8 | 1069.8 | 1069.8 | 1069.8 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.0000 |
| 0.2149 | 1121.7 | 1134.7 | 977.9 | 1116.3 | 1164.4 | 1101.5 | 1.15 | -14.70 | -0.48 | 3.66 | -1.83 | 1.0097 |
| 0.3811 | 1170.5 | 1188.8 | 911.1 | 1160.8 | 1229.5 | 1135.9 | 1.54 | -28.48 | -0.84 | 4.80 | -3.04 | 1.0168 |
| 0.5135 | 1213.5 | 1234.5 | 876.7 | 1202.6 | 1277.1 | 1171.6 | 1.70 | -38.42 | -0.90 | 4.98 | -3.57 | 1.0182 |
| 0.6215 | 1253.1 | 1273.7 | 862.8 | 1241.7 | 1313.3 | 1208.0 | 1.61 | -45.24 | -0.92 | 4.59 | -3.74 | 1.0185 |
| 0.7112 | 1290.8 | 1307.5 | 863.8 | 1278.0 | 1341.9 | 1244.5 | 1.28 | -49.42 | -1.00 | 3.81 | -3.72 | 1.0201 |
| 0.7870 | 1324.6 | 1337.1 | 886.0 | 1311.8 | 1365.0 | 1281.1 | 0.93 | -49.51 | -0.98 | 2.96 | -3.40 | 1.0196 |
| 0.8518 | 1355.1 | 1363.1 | 933.8 | 1343.2 | 1384.1 | 1317.5 | 0.59 | -45.11 | -0.88 | 2.09 | -2.85 | 1.0177 |
| 0.9079 | 1381.2 | 1386.2 | 1024.7 | 1372.5 | 1400.1 | 1353.7 | 0.36 | -34.79 | -0.63 | 1.35 | -2.03 | 1.0127 |
| 0.9568 | 1404.5 | 1406.9 | 1177.3 | 1399.9 | 1413.7 | 1389.7 | 0.17 | -19.30 | -0.33 | 0.65 | -1.06 | 1.0066 |
| 1.0000 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.0000 |
| 313K | | | | | | | | | | | | |
| 0.0000 | 1047.0 | 1047.0 | 1047.0 | 1047.0 | 1047.0 | 1047.0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.0000 |
| 0.2149 | 1095.7 | 1106.4 | 958.6 | 1091.6 | 1141.2 | 1071.4 | 0.97 | -14.30 | -0.37 | 3.99 | -2.27 | 1.0075 |
| 0.3811 | 1140.6 | 1156.8 | 902.3 | 1134.3 | 1204.5 | 1100.3 | 1.40 | -26.41 | -0.56 | 5.30 | -3.66 | 1.0112 |
| 0.5135 | 1181.2 | 1200.1 | 874.7 | 1174.3 | 1249.9 | 1132.0 | 1.57 | -35.03 | -0.58 | 5.49 | -4.34 | 1.0117 |
| 0.6215 | 1218.2 | 1237.5 | 867.7 | 1211.7 | 1284.1 | 1165.6 | 1.56 | -40.40 | -0.53 | 5.13 | -4.51 | 1.0107 |
| 0.7112 | 1254.6 | 1270.3 | 869.8 | 1246.5 | 1310.7 | 1200.6 | 1.24 | -44.24 | -0.65 | 4.28 | -4.50 | 1.0131 |
| 0.7870 | 1287.6 | 1299.2 | 892.4 | 1278.8 | 1332.1 | 1236.6 | 0.89 | -44.29 | -0.69 | 3.34 | -4.13 | 1.0138 |
| 0.8518 | 1317.0 | 1324.9 | 941.0 | 1308.8 | 1349.6 | 1273.3 | 0.60 | -39.96 | -0.62 | 2.41 | -3.43 | 1.0125 |
| 0.9079 | 1344.2 | 1347.9 | 1019.3 | 1336.8 | 1364.2 | 1310.7 | 0.27 | -31.88 | -0.55 | 1.47 | -2.55 | 1.0111 |
| 0.9568 | 1366.5 | 1368.5 | 1162.7 | 1362.9 | 1376.6 | 1348.7 | 0.15 | -17.52 | -0.27 | 0.73 | -1.32 | 1.0053 |
| 1.0000 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 1.0000 |
| 318K | | | | | | | | | | | | |
| 0.0000 | 1029.4 | 1029.4 | 1029.4 | 1029.4 | 1029.4 | 1029.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.2149 | 1077.2 | 1087.2 | 942.0 | 1074.0 | 1126.3 | 1050.0 | 0.9 | -14.4 | -0.3 | 4.4 | -2.6 | 1.0059 |
| 0.3811 | 1121.2 | 1136.7 | 887.9 | 1116.7 | 1190.2 | 1076.2 | 1.4 | -26.3 | -0.4 | 5.8 | -4.2 | 1.0081 |
| 0.5135 | 1161.6 | 1179.7 | 861.0 | 1156.8 | 1235.6 | 1106.2 | 1.5 | -34.9 | -0.4 | 6.0 | -5.0 | 1.0083 |
| 0.6215 | 1199.1 | 1217.3 | 853.1 | 1194.3 | 1269.5 | 1138.9 | 1.5 | -40.6 | -0.4 | 5.5 | -5.3 | 1.0081 |
| 0.7112 | 1235.0 | 1250.4 | 857.0 | 1229.1 | 1295.8 | 1173.7 | 1.2 | -44.1 | -0.5 | 4.7 | -5.2 | 1.0096 |
| 0.7870 | 1268.7 | 1279.8 | 876.5 | 1261.5 | 1316.8 | 1210.2 | 0.9 | -44.7 | -0.6 | 3.6 | -4.8 | 1.0114 |
| 0.8518 | 1299.5 | 1306.1 | 921.3 | 1291.7 | 1333.9 | 1248.3 | 0.5 | -41.1 | -0.6 | 2.6 | -4.1 | 1.0122 |
| 0.9079 | 1327.5 | 1329.7 | 994.1 | 1319.7 | 1348.1 | 1287.7 | 0.2 | -33.5 | -0.6 | 1.5 | -3.1 | 1.0118 |
| 0.9568 | 1350.5 | 1351.0 | 1125.0 | 1345.9 | 1360.1 | 1328.4 | 0.0 | -20.0 | -0.3 | 0.7 | -1.7 | 1.0068 |
| 1.0000 | 1370.4 | 1370.4 | 1370.4 | 1370.4 | 1370.4 | 1370.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |

Table 2. Experimental and Theoretical values of Ultrasonic velocity and Percentage of Deviation at different temperatures for the binary liquid mixture- II (NN-Dimethylacetamide + Diethanolamine).

| Mole Fraction of NNDA X_1 | Exp. Velocity U_{exp} | Theoretical Velocities | | | | | Percentage Deviation | | | | | $\frac{U^2}{U_{imx}^2}$ |
|--------------------------------|----------------------------|------------------------|-----------|-----------|-----------|------------|----------------------|-----------|-----------|-----------|------------|-------------------------|
| | | U_{Nom} | U_{FLT} | U_{IMR} | U_{IDR} | U_{JUNJ} | U_{Nom} | U_{FLT} | U_{IMR} | U_{IDR} | U_{JUNJ} | |
| 303K | | | | | | | | | | | | |
| 0.0000 | 1719.2 | 1719.2 | 1719.2 | 1719.2 | 1719.2 | 1719.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1690.8 | 1685.1 | 1984.9 | 1667.0 | 1689.9 | 1670.7 | -0.3 | 14.8 | -1.4 | -0.1 | -1.2 | 1.0287 |
| 0.2316 | 1663.0 | 1652.4 | 2678.3 | 1622.9 | 1660.7 | 1629.1 | -0.6 | 37.9 | -2.5 | -0.1 | -2.1 | 1.0501 |
| 0.3407 | 1635.6 | 1621.2 | 1097.6 | 1585.3 | 1631.8 | 1593.0 | -0.9 | 85.1 | -3.2 | -0.2 | -2.7 | 1.0645 |
| 0.4456 | 1608.0 | 1591.3 | -747.9 | 1553.1 | 1603.1 | 1561.6 | -1.1 | 315.0 | -3.5 | -0.3 | -3.0 | 1.0719 |
| 0.5466 | 1578.7 | 1562.6 | 409.4 | 1525.5 | 1574.7 | 1533.9 | -1.0 | -285.6 | -3.5 | -0.3 | -2.9 | 1.0709 |
| 0.6440 | 1548.7 | 1535.1 | 789.4 | 1501.8 | 1546.4 | 1509.5 | -0.9 | -96.2 | -3.1 | -0.1 | -2.6 | 1.0634 |
| 0.7378 | 1519.2 | 1508.7 | 1007.2 | 1481.3 | 1518.4 | 1487.8 | -0.7 | -50.8 | -2.6 | -0.1 | -2.1 | 1.0518 |
| 0.8283 | 1490.4 | 1483.3 | 1178.2 | 1463.7 | 1490.5 | 1468.4 | -0.5 | -26.5 | -1.8 | 0.0 | -1.5 | 1.0368 |
| 0.9156 | 1462.5 | 1459.0 | 1314.9 | 1448.5 | 1462.9 | 1451.1 | -0.2 | -11.2 | -1.0 | 0.0 | -0.8 | 1.0194 |
| 1.0000 | 1435.5 | 1435.5 | 1435.5 | 1435.5 | 1435.5 | 1435.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 308K | | | | | | | | | | | | |
| 0.0000 | 1698.8 | 1698.8 | 1698.8 | 1698.8 | 1698.8 | 1698.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1672.4 | 1665.6 | 2085.3 | 1648.5 | 1670.9 | 1650.9 | -0.4 | 19.8 | -1.4 | -0.1 | -1.3 | 1.0292 |
| 0.2316 | 1646.0 | 1633.9 | 3658.0 | 1606.0 | 1643.1 | 1610.1 | -0.7 | 55.0 | -2.5 | -0.2 | -2.2 | 1.0505 |
| 0.3407 | 1620.0 | 1603.6 | 2729.0 | 1569.7 | 1615.4 | 1575.0 | -1.0 | 159.4 | -3.2 | -0.3 | -2.9 | 1.0651 |
| 0.4456 | 1593.4 | 1574.7 | 153.1 | 1538.7 | 1587.8 | 1544.6 | -1.2 | -940.9 | -3.6 | -0.3 | -3.2 | 1.0723 |
| 0.5466 | 1565.5 | 1547.1 | 649.5 | 1512.1 | 1560.4 | 1518.1 | -1.2 | -141.0 | -3.5 | -0.3 | -3.1 | 1.0719 |
| 0.6440 | 1536.5 | 1520.6 | 884.5 | 1489.2 | 1533.2 | 1494.8 | -1.0 | -73.7 | -3.2 | -0.2 | -2.8 | 1.0645 |
| 0.7378 | 1507.5 | 1495.3 | 1046.4 | 1469.5 | 1506.0 | 1474.2 | -0.8 | -44.1 | -2.6 | -0.1 | -2.3 | 1.0524 |
| 0.8283 | 1479.5 | 1471.0 | 1185.9 | 1452.5 | 1479.0 | 1456.0 | -0.6 | -24.8 | -1.9 | 0.0 | -1.6 | 1.0375 |
| 0.9156 | 1452.4 | 1447.8 | 1302.2 | 1437.9 | 1452.1 | 1439.8 | -0.3 | -11.5 | -1.0 | 0.0 | -0.9 | 1.0202 |
| 1.0000 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 313K | | | | | | | | | | | | |
| 0.0000 | 1687.0 | 1687.0 | 1687.0 | 1687.0 | 1687.0 | 1687.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1658.1 | 1650.2 | 2319.0 | 1631.7 | 1656.5 | 1633.0 | -0.5 | 28.5 | -1.6 | -0.1 | -1.5 | 1.0326 |
| 0.2316 | 1628.8 | 1615.3 | 2659.7 | 1585.0 | 1626.0 | 1587.6 | -0.8 | 93.9 | -2.8 | -0.2 | -2.6 | 1.0560 |
| 0.3407 | 1600.4 | 1581.9 | 180.1 | 1545.4 | 1595.7 | 1548.8 | -1.2 | 988.6 | -3.6 | -0.3 | -3.3 | 1.0725 |
| 0.4456 | 1571.2 | 1550.2 | 548.3 | 1511.5 | 1565.6 | 1515.6 | -1.4 | -186.5 | -3.9 | -0.4 | -3.7 | 1.0806 |
| 0.5466 | 1540.6 | 1519.9 | 806.9 | 1482.4 | 1535.5 | 1486.7 | -1.4 | -90.9 | -3.9 | -0.3 | -3.6 | 1.0800 |
| 0.6440 | 1509.2 | 1490.9 | 959.6 | 1457.4 | 1505.6 | 1461.5 | -1.2 | -57.3 | -3.6 | -0.2 | -3.3 | 1.0724 |
| 0.7378 | 1478.0 | 1463.3 | 1075.9 | 1435.8 | 1475.8 | 1439.4 | -1.0 | -37.4 | -2.9 | -0.1 | -2.7 | 1.0596 |
| 0.8283 | 1447.2 | 1436.8 | 1184.1 | 1417.2 | 1446.2 | 1419.9 | -0.7 | -22.2 | -2.1 | -0.1 | -1.9 | 1.0428 |
| 0.9156 | 1417.2 | 1411.5 | 1279.4 | 1401.1 | 1416.6 | 1402.6 | -0.4 | -10.8 | -1.2 | 0.0 | -1.0 | 1.0232 |
| 1.0000 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 318K | | | | | | | | | | | | |
| 0.0000 | 1667.4 | 1667.4 | 1667.4 | 1667.4 | 1667.4 | 1667.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1639.2 | 1631.1 | 3003.4 | 1613.0 | 1637.5 | 1613.8 | -0.5 | 45.4 | -1.6 | -0.1 | -1.6 | 1.0327 |
| 0.2316 | 1611.2 | 1596.6 | 1155.6 | 1567.1 | 1607.6 | 1568.8 | -0.9 | 239.4 | -2.8 | -0.2 | -2.7 | 1.0571 |
| 0.3407 | 1583.6 | 1563.7 | 403.1 | 1528.1 | 1577.9 | 1530.6 | -1.3 | -292.8 | -3.6 | -0.4 | -3.5 | 1.0740 |
| 0.4456 | 1555.2 | 1532.5 | 728.6 | 1494.7 | 1548.2 | 1497.8 | -1.5 | -113.4 | -4.0 | -0.4 | -3.8 | 1.0826 |
| 0.5466 | 1525.2 | 1502.7 | 888.1 | 1466.1 | 1518.7 | 1469.5 | -1.5 | -71.7 | -4.0 | -0.4 | -3.8 | 1.0823 |
| 0.6440 | 1494.2 | 1474.2 | 999.1 | 1441.5 | 1489.2 | 1444.8 | -1.4 | -49.6 | -3.7 | -0.3 | -3.4 | 1.0745 |
| 0.7378 | 1463.1 | 1447.0 | 1094.0 | 1420.2 | 1459.9 | 1423.2 | -1.1 | -33.7 | -3.0 | -0.2 | -2.8 | 1.0613 |
| 0.8283 | 1432.3 | 1421.1 | 1186.7 | 1401.9 | 1430.6 | 1404.1 | -0.8 | -20.7 | -2.2 | -0.1 | -2.0 | 1.0439 |
| 0.9156 | 1402.2 | 1396.2 | 1275.7 | 1386.0 | 1401.5 | 1387.3 | -0.4 | -9.9 | -1.2 | -0.1 | -1.1 | 1.0234 |
| 1.0000 | 1372.4 | 1372.4 | 1372.4 | 1372.4 | 1372.4 | 1372.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |

Table 3. Experimental and Theoretical values of Ultrasonic velocity and Percentage of Deviation at different temperatures for the binary liquid mixture- II (NN-Dimethylacetamide + Benzylamine).

| Mole Fraction of NNDA X_1 | Exp. Velocity U_{exp} | Theoretical Velocities | | | | | Percentage Deviation | | | | | $\frac{U^2}{U_{imx}^2}$ |
|--------------------------------|----------------------------|------------------------|-----------|-----------|-----------|------------|----------------------|-----------|-----------|-----------|------------|-------------------------|
| | | U_{Nom} | U_{FLT} | U_{IMR} | U_{IDR} | U_{JUNJ} | U_{Nom} | U_{FLT} | U_{IMR} | U_{IDR} | U_{JUNJ} | |
| 303K | | | | | | | | | | | | |
| 0.0000 | 1719.2 | 1719.2 | 1719.2 | 1719.2 | 1719.2 | 1719.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1690.8 | 1685.1 | 1984.9 | 1667.0 | 1689.9 | 1670.7 | -0.3 | 14.8 | -1.4 | -0.1 | -1.2 | 1.0287 |
| 0.2316 | 1663.0 | 1652.4 | 2678.3 | 1622.9 | 1660.7 | 1629.1 | -0.6 | 37.9 | -2.5 | -0.1 | -2.1 | 1.0501 |
| 0.3407 | 1635.6 | 1621.2 | 1097.6 | 1585.3 | 1631.8 | 1593.0 | -0.9 | 85.1 | -3.2 | -0.2 | -2.7 | 1.0645 |
| 0.4456 | 1608.0 | 1591.3 | -747.9 | 1553.1 | 1603.1 | 1561.6 | -1.1 | 315.0 | -3.5 | -0.3 | -3.0 | 1.0719 |
| 0.5466 | 1578.7 | 1562.6 | 409.4 | 1525.5 | 1574.7 | 1533.9 | -1.0 | -285.6 | -3.5 | -0.3 | -2.9 | 1.0709 |
| 0.6440 | 1548.7 | 1535.1 | 789.4 | 1501.8 | 1546.4 | 1509.5 | -0.9 | -96.2 | -3.1 | -0.1 | -2.6 | 1.0634 |
| 0.7378 | 1519.2 | 1508.7 | 1007.2 | 1481.3 | 1518.4 | 1487.8 | -0.7 | -50.8 | -2.6 | -0.1 | -2.1 | 1.0518 |
| 0.8283 | 1490.4 | 1483.3 | 1178.2 | 1463.7 | 1490.5 | 1468.4 | -0.5 | -26.5 | -1.8 | 0.0 | -1.5 | 1.0368 |
| 0.9156 | 1462.5 | 1459.0 | 1314.9 | 1448.5 | 1462.9 | 1451.1 | -0.2 | -11.2 | -1.0 | 0.0 | -0.8 | 1.0194 |
| 1.0000 | 1435.5 | 1435.5 | 1435.5 | 1435.5 | 1435.5 | 1435.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 308K | | | | | | | | | | | | |
| 0.0000 | 1698.8 | 1698.8 | 1698.8 | 1698.8 | 1698.8 | 1698.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1672.4 | 1665.6 | 2085.3 | 1648.5 | 1670.9 | 1650.9 | -0.4 | 19.8 | -1.4 | -0.1 | -1.3 | 1.0292 |
| 0.2316 | 1646.0 | 1633.9 | 3658.0 | 1606.0 | 1643.1 | 1610.1 | -0.7 | 55.0 | -2.5 | -0.2 | -2.2 | 1.0505 |
| 0.3407 | 1620.0 | 1603.6 | 2729.0 | 1569.7 | 1615.4 | 1575.0 | -1.0 | 159.4 | -3.2 | -0.3 | -2.9 | 1.0651 |
| 0.4456 | 1593.4 | 1574.7 | 153.1 | 1538.7 | 1587.8 | 1544.6 | -1.2 | -940.9 | -3.6 | -0.3 | -3.2 | 1.0723 |
| 0.5466 | 1565.5 | 1547.1 | 649.5 | 1512.1 | 1560.4 | 1518.1 | -1.2 | -141.0 | -3.5 | -0.3 | -3.1 | 1.0719 |
| 0.6440 | 1536.5 | 1520.6 | 884.5 | 1489.2 | 1533.2 | 1494.8 | -1.0 | -73.7 | -3.2 | -0.2 | -2.8 | 1.0645 |
| 0.7378 | 1507.5 | 1495.3 | 1046.4 | 1469.5 | 1506.0 | 1474.2 | -0.8 | -44.1 | -2.6 | -0.1 | -2.3 | 1.0524 |
| 0.8283 | 1479.5 | 1471.0 | 1185.9 | 1452.5 | 1479.0 | 1456.0 | -0.6 | -24.8 | -1.9 | 0.0 | -1.6 | 1.0375 |
| 0.9156 | 1452.4 | 1447.8 | 1302.2 | 1437.9 | 1452.1 | 1439.8 | -0.3 | -11.5 | -1.0 | 0.0 | -0.9 | 1.0202 |
| 1.0000 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 1425.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 313K | | | | | | | | | | | | |
| 0.0000 | 1687.0 | 1687.0 | 1687.0 | 1687.0 | 1687.0 | 1687.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1658.1 | 1650.2 | 2319.0 | 1631.7 | 1656.5 | 1633.0 | -0.5 | 28.5 | -1.6 | -0.1 | -1.5 | 1.0326 |
| 0.2316 | 1628.8 | 1615.3 | 2659.7 | 1585.0 | 1626.0 | 1587.6 | -0.8 | 93.9 | -2.8 | -0.2 | -2.6 | 1.0560 |
| 0.3407 | 1600.4 | 1581.9 | 180.1 | 1545.4 | 1595.7 | 1548.8 | -1.2 | 988.6 | -3.6 | -0.3 | -3.3 | 1.0725 |
| 0.4456 | 1571.2 | 1550.2 | 548.3 | 1511.5 | 1565.6 | 1515.6 | -1.4 | -186.5 | -3.9 | -0.4 | -3.7 | 1.0806 |
| 0.5466 | 1540.6 | 1519.9 | 806.9 | 1482.4 | 1535.5 | 1486.7 | -1.4 | -90.9 | -3.9 | -0.3 | -3.6 | 1.0800 |
| 0.6440 | 1509.2 | 1490.9 | 959.6 | 1457.4 | 1505.6 | 1461.5 | -1.2 | -57.3 | -3.6 | -0.2 | -3.3 | 1.0724 |
| 0.7378 | 1478.0 | 1463.3 | 1075.9 | 1435.8 | 1475.8 | 1439.4 | -1.0 | -37.4 | -2.9 | -0.1 | -2.7 | 1.0596 |
| 0.8283 | 1447.2 | 1436.8 | 1184.1 | 1417.2 | 1446.2 | 1419.9 | -0.7 | -22.2 | -2.1 | -0.1 | -1.9 | 1.0428 |
| 0.9156 | 1417.2 | 1411.5 | 1279.4 | 1401.1 | 1416.6 | 1402.6 | -0.4 | -10.8 | -1.2 | 0.0 | -1.0 | 1.0232 |
| 1.0000 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 1387.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 318K | | | | | | | | | | | | |
| 0.0000 | 1667.4 | 1667.4 | 1667.4 | 1667.4 | 1667.4 | 1667.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |
| 0.1181 | 1639.2 | 1631.1 | 3003.4 | 1613.0 | 1637.5 | 1613.8 | -0.5 | 45.4 | -1.6 | -0.1 | -1.6 | 1.0327 |
| 0.2316 | 1611.2 | 1596.6 | 1155.6 | 1567.1 | 1607.6 | 1568.8 | -0.9 | 239.4 | -2.8 | -0.2 | -2.7 | 1.0571 |
| 0.3407 | 1583.6 | 1563.7 | 403.1 | 1528.1 | 1577.9 | 1530.6 | -1.3 | -292.8 | -3.6 | -0.4 | -3.5 | 1.0740 |
| 0.4456 | 1555.2 | 1532.5 | 728.6 | 1494.7 | 1548.2 | 1497.8 | -1.5 | -113.4 | -4.0 | -0.4 | -3.8 | 1.0826 |
| 0.5466 | 1525.2 | 1502.7 | 888.1 | 1466.1 | 1518.7 | 1469.5 | -1.5 | -71.7 | -4.0 | -0.4 | -3.8 | 1.0823 |
| 0.6440 | 1494.2 | 1474.2 | 999.1 | 1441.5 | 1489.2 | 1444.8 | -1.4 | -49.6 | -3.7 | -0.3 | -3.4 | 1.0745 |
| 0.7378 | 1463.1 | 1447.0 | 1094.0 | 1420.2 | 1459.9 | 1423.2 | -1.1 | -33.7 | -3.0 | -0.2 | -2.8 | 1.0613 |
| 0.8283 | 1432.3 | 1421.1 | 1186.7 | 1401.9 | 1430.6 | 1404.1 | -0.8 | -20.7 | -2.2 | -0.1 | -2.0 | 1.0439 |
| 0.9156 | 1402.2 | 1396.2 | 1275.7 | 1386.0 | 1401.5 | 1387.3 | -0.4 | -9.9 | -1.2 | -0.1 | -1.1 | 1.0234 |
| 1.0000 | 1372.4 | 1372.4 | 1372.4 | 1372.4 | 1372.4 | 1372.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0000 |

V. CONCLUSION

In the binary liquid mixtures, N,N Dimethyl acetamide + Diethylamine, N,N Dimethyl acetamide + Diethanolamine, N,N Dimethyl acetamide+ Benzylamine it is observed that there is a close agreement between experimental and theoretical values calculated by Nomoto's relation, followed by Junjie's, Ideal Mixing Relation and Impedance Dependence Relation. It may be concluded that out of the five theories and relations discussed above, the Nomoto's relation, Junjie's relation, Ideal Mixing Relation and Impedance Dependence Relation are good for estimation of velocities, as the values obtained show good agreement with the respective measured values of ultrasonic velocity.

REFERENCES

- [1] U. Sridevi, K. Samatha and A. Visvanthasarma, (2004), Excess thermodynamic properties in binary liquids, J. Pure Appl. Ultrasonics, 26, pp. 1-11.
- [2] P.P.Divakar, B.V.Rao, K.Samatha ,(2012),Study of Molecular Interactions in Binary Liquid Mixtures by Acoustical Method at 303K, E-Journal of Chemistry , 9(3),pp.1332-1335
- [3] A N Kannappan and R palani, (2007), ultrasonic investigations in amino acids with aqueous dimethyl formamide, Indian J. Chem ,Indian J. Chem , 54, 46A.
- [4] P.P.Divakar, K.Samatha,(2015),comparative Study of Experimental and Theoretical Ultrasonic Velocities in Binary Mixtures of Cyclohexanone with Aliphatic Esters at Different Temperatures, International Letters of Chemistry, Physics and Astronomy 3,pp. 13-24
- [5] J. N. Ramteke and S. B. Khasare,(2012),Comparison of Theoretical Ultrasonic Velocities in Binary Liquid Mixtures Containing a-Picoline in Ethanol Advances in Applied Science Research, 3 (6),pp.3415-3420.
- [6] Vaidya Rohit, K.S., N.K., M.(2015) : Theoretical models of ultrasonic velocities in binary liquid mixtures. ResearchJournal of Chemical Sciences 5(10), 33
- [7] O.Nomoto,(1958),Empirical formula for sound velocities in liquid mixtures, Journal of the Physical Society of Japan,13,pp.1528-1532.
- [8] Jacobson B, ActaChemScand, (1952),6, pp. 1485, & J. Chem. Phys, ,(1952), 20,pp.927.
- [9] W.VanDael and Vangeel, in Proceedings of the International Conference on Calorimetry and Thermodynamics, (1955) pp.555, Warsaw, Poland.
- [10] Bladamer M and Waddington D, (1970), J Phys Chem, 74, pp.2569.
- [11] W. Schaaffs(1939), Z. phys, 114,pp. 110., &Schaaffs W (1963) , Molekularakustik, Springer-Verlag, Berlin.
- [12] Junjie Z, (1984),J China UnivSci Tech., 14, pp.298.
- [13] Rao GVR, Samantha K, Sarma AV (2004) A comparative study of ultrasonic velocity and allied parameters of binary mixtures at different temperatures. J.AcoustSocInd 32, pp.213.
- [14] Dubey GP, Monika Sharma (2008) Acoustical and excess properties of {1-hexanol + n-hexane, or n-octane, or n-decane} at (298.15, 303.15, and 308.15) K. Journal of Molecular Liquids, 142, pp. 124-129.
- [15] Artigas HI Bandrés, B Giner, C & A.L afuenteVillares, (2008), J. MolLiq, 139, pp.138.
- [16] Radhamma M, VenkatesuP, Prabhakara Rao MV, Lee MJ, Lin HM (2008), Excess molar volumes and ultrasonic studies of dimethyl sulphoxide with ketones at T = 303.15 K. The Journal of Chemical Thermodynamics, 40, pp. 492-497.
- [17] FakruddinSk, SrinivasuCh, Narendra K (2012), Theoretical Studies of Ultrasonic Velocities in Binary Liquid Mixtures of Quinoline at Different Temperatures. Journal of Chemical and Pharmaceutical Research 4, pp. 1799-1806.
- [18] Rita Mehra, Avneesh K Gaur (2008) Study of a Binary Liquid Mixture of Diethylamine and 1-Decanol and Validation of Theoretical Approaches of Sound Speed at Different Temperatures. J ChemEng Data 53, pp. 863-866.
- [19] Gliński(2002), Additivity of sound velocity in liquid mixtures, J.Sol. Chemistry,31,pp. 59.
- [20] Suhashini Ernest, Kavitha P (2011) Theoretical evaluation of Ultrasonic velocity in Binary mixtures of an edible oil with alkyl Acetates. International Journal ofChemical, Environmental and Pharmaceutical Research 2, pp. 92-95.
- [21] Santhi, N., Madumitha, (2014) J.: Molecular interaction studies in binary liquid mixture through ultrasonic measurements at 303.15k. International Journal of Advanced Chemistry 4(1), 227.
- [22] Shukla B D, Jha L K and Dwivedi D K,(1991), J Pure Appl, Ultrason, 13, pp. 72.

Authors



Vysyaraju Sravanthi, has been working as a PGT in AP Model School, Vizianagaram and pursuing her Ph.D (part-Time) under the guidance of Prof. K. Samatha, Dept. of Physics, Andhra University, Visakhapatnam, India.



Dr. Putla Paul Divakar has been working as associate professor in Dept. of Physics, Sir C. R. Reddy Autonomous College, Eluru, India. The publications are 14, so far in various international and national journals. He is General Assembly Member of AIRAPT, France and also Life Member of Indian Science Congress. His profile was incorporated in Marqui's Who's Who-2014, USA.



Prof. K. Samatha, has been working as a Professor in the Dept. of Physics, Andhra University, Visakhapatnam, India. She has been working in the fields of Solid State Physics and Ultrasonics. She has published 45 papers so far in various national / international journals. Till now under her guidance 10 Ph. D Scholars and 08 M .Phil scholars were awarded with respective degrees.